

COMPLEMENTARY THERAPY IN POLYCYSTIC OVARY SYNDROME

Aquino C. I., Nori S. L.

Department of Medicine and Surgery University of Salerno, Via S Allende, Baronissi
Corresponding author: Carmen Imma Aquino (C.AQUINO8@STUDENTI.UNISA.IT)

Abstract – Polycystic Ovary Syndrome (PCOS) is an endocrine disease. PCOS afflicts 5 to 10 % of women of reproductive age. The symptoms are: amenorrhea, oligomenorrhea, hirsutism, obesity, infertility, chronic hyperandrogenic anovulation and acne.

Other risk factors aggravate this condition: insulin resistance, obesity, hypertension, dyslipidemia, inflammation and subclinical cardiovascular disease. Anxiety, depression and reduced quality of life are also common.

This review highlights the mechanisms and the beneficial effects of acupuncture, exercise and resveratrol on animal models and on humans affected by PCOS.

Keywords: PCOS, acupuncture, exercise, resveratrol.

I. INTRODUCTION

Polycystic Ovary Syndrome (PCOS) is an endocrine disease characterized by a still unclear aetiology resulting from the interaction among genetic, behavioral and environmental conditions and where ovarian hyperandrogenemia plays a key role [1].

Indeed, several studies demonstrate that a low dose of testosterone in female monkeys Rhesus increases luteinizing hormone (LH) secretion [2, 3, 4, 5]. Further, androgens can determine PCOS-like symptomatic pictures in animal female and female-to-male transsexual models [6].

Hyperinsulinemia/insulin resistance and abdominal obesity are relevant. However, whether hyperandrogenism results from the hyperinsulinemia of insulin resistance or vice versa is unclear [7, 8]. Studies performed on monkeys, sheep and rats demonstrate that prenatal androgenization is a cause of PCOS [9, 10]. In human, only one study has verified the presence of testosterone in the blood of babies from PCOS mothers [11]. Also the exposition to androgens in a prepuberal period causes PCOS in animal models [12, 13]. At the begin of ovary development, antimüllerian hormone (AMH) is a determinant of follicular activity. In rats increased AMH, as in PCOS, decreases the follicular growth in ovary [14, 15] and is linked with low levels of FSH [16, 17]. Moreover, as shown from in vitro studies on granulosa

cells of small bovine follicles cultured with testosterone, AMH expression is downregulated and may cause PCOS [18].

It is known that neuroendocrine defects can increase the persistently rapid LH pulsatility- and this latter have been linked with augment of the ovarian volume, several antral follicles and androgen hyperproduction [8]. The high activity in the sympathetic neurons innervating the ovaries precedes the development of ovarian cysts in rats [16]. Women with PCOS may have increased ovarian nerve fiber density and androgen hypersecretion. Further, the high circulating testosterone in PCOS women explains their high sympathetic nervous system activity [17] associated with several clinical findings illustrated in Table 1 [19].

TABLE 1. SIGNS AND SYMPTOMS OF PCOS CAUSED BY INCREASED ACTIVITY IN SNS

Hyperandrogenism
Increased sympathetic nerve activity
Altered GH/IGF-1 axis
Hypersecretion of LH
Anovulation
Visceral obesity
Hypertension
Insulin resistance
Psychological stress

LH, Luteinising hormone;
GH, growth hormone; IGF, insulin growth factor.

Reprinted with modifications from E. Stener-Victorin et al. (2008) with permission from John Wiley and Sons, copyright (2013) License Number 3286560319536

It has been demonstrated that PCOS women have ovarian theca interna hyperplasia responsible for the increased androgen steroidogenesis. Besides, theca cells have higher expression of LH receptors with augmented sensibility to LH stimulation [18, 20, 21].

GnRH pulse generator is inhibited by the central β -endorphin system that also modulates sympathetic tone. Diminished or augmented β -endorphin and dopamine

activity causes tonic increasing release of LH [22, 23]. High plasma β -endorphin levels are related to hyperinsulinemia, stress [24, 25] and also to PCOS. Growth hormone (GH) and insulin growth factor-1 (IGF1) regulate sympathetic nerve tone : and in PCOS have been shown to be present disturbances in the somatotrophic axis (GH/ IGF-1) [26]. Moreover, high sympathetic nervous tone is the most relevant risk factor for heart diseases and mortality [27, 28]. The effects of these alterations are shown in the figure 1 [29, 30, 4].




Fig. 1: Schematic illustration of PCOS events. Aquino C. I. , 2013

II. METHODOLOGY

1. ACUPUNCTURE

Because PCOS is a multifactorial pathology several treatments (see table 2) have been proposed [31]. It is demonstrated that cognitive performances (word learning test and symbol digit substitution test) are also improved significantly after the treatment (Metformin) for PCOS [32]. Acupuncture is now widely used in the Western population accounting for one in three Americans using it [5].

TABLE 2. TREATMENTS OF PCOS

- OLD THERAPY
- Oral contraceptives (OC)
- Anti-androgens
- NEW THERAPY
- Insulin sensitizers:
- (Metformin Thiazolidinediones)
- EMERGING THERAPY
- Statins
- Acupuncture
- Dietary products and nutrients.
- Vitamin D
- Herbal medicines
- Vitamin B12 and folate.
- Resveratrol
- AGEs low diet
- Physical activity

Acupuncture is an ancient beneficial practice that use needles for manual or electrical sensory stimulation of somatic afferent nerves. These nerves innervate the skin and muscles.

Auricular stimulation is widely used to improve pregnancy (see table of auricular acupuncture in figure 2). Traditional Chinese medicine (TCM) affirmed that acupuncture restores the balance between Yin and Yang. In the modern interpretation of the Western Medicine, acupuncture restores the equilibrium between parasympathetic and sympathetic activity [33].

The insertion of needles determines a specific pattern of afferent response in peripheral nerves stimulating the spinal cord and the central nervous system [34].




Fig.2: Auricular acupuncture. Reprinted from Ernest Hung Yu Ng et al. (2008), by permission from Elsevier, copyright (2013) License number 3286611228365.

Electroacupuncture (EA) takes advantage of a low frequency (1-15 Hz) electrical stimulation that elicits

muscle contraction and augments the analgesic effects of acupuncture [35]. Acupuncture can activate autonomic responses (as seen in figure 3), the descending inhibitory pain control systems, and the propriospinal heterosegmental antinociceptive C-fibers [36, 37] and α , β , δ fibers [38, 34]. Positive results of this millenary treatment for PCOS women is related to: the central sympathetic inhibition through a major release of endorphin, the change of uterine blood flow and motility, the stress reduction. Acupuncture regulates the hypothalamic-pituitary-ovarian axis (HPO) actions by modulating central opioids, in particular β -endorphin [38, 39] in a context of mesencephalic and brainstem network [40, 41].




Fig. 4: localization of ST36 in the tibialis anterior muscle, 4 fingerbreadths below the kneecap and 1 fingerbreadth lateral from the anterior crest of the tibia [54]. Reprinted from Kathleen K.S. Hui et al. (2005), by permission from Elsevier, copyright (2013) License number 3286610770603




Fig. 3: EA and autonomic effects. Reprinted with modifications from E. Stener-Victorin et al. (2010), with permission of Elsevier. copyright (2013), License number 3286600811085

There are opioid receptors of several types: μ , δ and κ in peripheral afferent nerve terminals [42, 43]. β -endorphin is derived from proopiomelanocortin (POMC). It is produced in the arcuate nucleus of hypothalamus and in the pituitary gland and binds μ receptors [44, 45, 46]. This nervous network is composed by: hypothalamus, medulla oblongata, ventrolateral periaqueductal gray, and the dorsomedial prefrontal cortex. This functional circuit appear to be very important for the autonomic regulation [47, 48], in particular for the pain modulation. Further, functional MRI and PET studies demonstrate that using acupuncture is possible to show significant modulatory effects on the limbic system, paralimbic and subcortical gray structures [49, 50, 51, 52, 53, 38]. An important acupuncture point, used also in PCOS and in other disorders, is ST 36 (stomach 36 or Zusanli; shown in figure 4) on the lower leg .

When it is stimulated, fRMN shows a predominance of signal attenuation, particularly in the limbic/paralimbic regions of the telencephalon, diencephalon and brainstem [50, 51, 53].

This evidences an integrated response of the human cerebro-cerebellar and limbic systems to acupuncture treatment. As shown also in rats models, after acupuncture, fRMN and PET results indicate an integrated response from multiple functional units of the brain which appear to be dependent on the psychophysical status . Moreover, the hemodynamic response correlates with acupuncture effects on the monoaminergic systems. Maybe these neuromodulators, in particular dopamine, may play an important role in acupuncture actions [54].

Another important beneficial effect of acupuncture is related to the actions on the corticotrophin releasing hormone (CRH)-produced by the hypothalamus. β -endorphin and reproductive functions are linked to GnRH and LH release.

Estrogens are important to ovulate for the LH surge [55] and opioids inhibits the effect of estrogens [56]. CRH is released under stress conditions and decreases GnRH secretion. Acupuncture decreases the levels of CRH and modulate the HPO and HPA axis [57].

Moreover, acupuncture have been shown to increase glucose uptake and microcirculation [58, 59].

Needles stimulations decrease sympathetic activity and increase parasympathetic activity inhibiting the dorsomedial prefrontal cortex [60]. Peripheral nerves release neuropeptides: neuropeptide Y (NPY), vasoactive intestinal polypeptide (VIP), substance P and calcitonin gene-related peptide (CGRP) with a rapid reaction [61, 62].

EA stimulates enkephalinergic neurons in the brain areas that regulates sympathetic outflow [63, 64].

According to these results, Li et al. [65] found that a single brief acupuncture stimulation at points P5-P6 transiently induces the production of enkephalin associated with a brain regulation of the sympathetic outflow.

Consequently, acupuncture may control the hypothalamic-pituitary-adrenal (HPA) axis (diminishing cortisol concentrations) and the hypothalamic-pituitary-gonadal (HPG) axis (modulating central β -endorphin production and secretion) [66]. EA controls the sympathetic reflex responses by decreasing the release of GABA in the ventrolateral periaqueductal gray (vlPAG) [67]. The decrease of GABA disinhibits vlPAG cells. This action modulates the activity of rostral ventrolateral medulla (rVLM) neurons attenuating the sympathoexcitatory reflex responses [68]. Probably, both glutamate and GABA are involved in the mechanism of acupuncture modulation of autonomic response.

Moreover, EA could give benefits to PCOS and other metabolic disorders, by altering sympathetic signaling [69, 70]. EA reduces mRNA^{NGF} in dihydrotestosterone (DHT)-induced rats and also in estradiol-valerate – (EV-) induced rats which have been used as PCOS models [71]. Nerve growth factor (NGF) is, in fact, higher in PCOS women ovaries and in EV-induced rats PCOS models [72, 73, 74]. NGF arrests follicle growth, augments ovarian sensibility to gonadotropins, increases adrenergic receptors and it causes ovarian hyperinnervation in transgenic mice models generated by pronuclear microinjection of the transgene construct into fertilized eggs from B6D2F1/J mice followed by transfer of the injected eggs into the uterus of a surrogate mother [72].

The decrease of NGF expression in peripheral organs could improve EA by modulating the autonomic nervous system activity [75]. In rats, EA decreases circulating testosterone concentrations and modulate the expression of opioid receptors μ and κ in the hypothalamus [76].

Furthermore, acupuncture in rats normalizes estrous cyclicity [76, 77], improves ovarian morphology [70], reduces circulating concentrations of inhibin B [78] and increases progesterone levels [76].

2. PHYSICAL ACTIVITY AND RESVERATROL

As shown in the figure 1, PCOS is characterized by lipidic blood profile, bigger adipocytes—and insulin resistance. These alterations determine visceral obesity and hypertension [79, 80]. Insulin resistance in skeletal muscles is related to alterations in fatty acid metabolism and mitochondrial abnormalities. In PCOS women it is associated with reduced gene expression of peroxisome proliferator-activated receptor γ coactivator α (PGC-1 α) and sirtuins (SIRT) 1 and 3 [81, 82].

Resveratrol, a natural polyphenol, may have positive metabolic effects. It balances lipid profile, decreases adiposity and improves insulin sensitivity [83]. Resveratrol increases, in skeletal muscle, the mRNA levels of estrogen-related receptor α (*Erra*), PGC-1 α , and the ERR α /PGC-1 α signaling pathway target gene

expression of nuclear respiratory factor 1 (*Nrf1*) (all decreased in PCOS) [84, 85].

Resveratrol causes apoptosis and diminishes proliferation of rat theca cells, reduces androgen levels by arresting Cyp17a1 gene expression [86, 87]. Some of these results are linked to the activation of SIRT 1 [88]. Resveratrol can activate AMP-activated protein kinase (AMPK) using a cAMP signaling and mediates glucose transporter (GLUT)-4 translocation and higher glucose uptake [89]. High leptin levels inhibit ovulation in rats [90] : this polyphenol combined to physical activity reduce fat mass and also leptin [69, 91]. Moreover, physical activity and somministration of resveratrol improve estrus cyclicity in PCOS rats [92]. The study by A. Benrick et al. by treating PCOS rats with resveratrol demonstrates that resveratrol has beneficial effects on adiposity (that improves) and cyclicity in a similar manner to exercise, but it is not a good candidate for treating insulin resistance associated with PCOS [92].

Moreover, physical activity and acupuncture decrease high muscle sympathetic nerve activity in women affected by PCOS. It is also demonstrated that acupuncture is more effective than physical exercises [18,93]. Physical activities determine weight loss improving cardiovascular order and insulin cellular response; visceral adiposity and insulin response (IR) decrease [94, 95].

Regular and moderate aerobic exercises (90 min per week at 60-70 % VO2max) over a short period improve ovulation and menstrual frequency [96, 97].

III.RESULTS

Goal of these review is to make a transactional correlation between the results obtained in PCOS animal models and human using acupuncture, physical activities and resveratrol. In the tables 3, 4 and 5 some of the experiments performed in this field are compared.

TABLE 3: EFFECTS OF ACUPUNCURE ON RODENTS MODELS

References	Description	Results
1 L. Manni, et al. 2005.	Virgin adult Wistar Kyoto rats were divided into four experimental groups: i) an Oil group (control, n = 8), ii) an Oil group receiving EA (EA, n = 8), iii) a PCO group (PCO, n = 8), and iv) a PCO group receiving EA (PCO+EA, n = 8). Daily stimulation were bilateral in the mm. biceps femoris and erector spinae, in somatic segments corresponding to the innervation of the ovaries.	EA modulated positively mRNA of PCOS rat models
2 J. Johansson et al., 2013	In rats with DHT-induced PCOS intense (5 days/wk) low-frequency EA for 4-5 wks.	Manual stimulation of acupuncture needles has a greater effect on glucose tolerance
3 E. Stener-Victorin et al., 2000	12 treatments by analyzing NGF in the central nervous system and the endocrine organs. Higher concentrations of NGF were found in the ovaries and the adrenal glands in the rats than in the control rats.	EA inhibits hyperactivity in the sympathetic nervous system.

TABLE 4: EFFECTS OF ACUPUNCTURE ON HUMANS WITH PCOS

References	Description	Results
4 J. Johansson et al., 2013	Manual acupuncture with EA (2Hz) using needles placed in abdominal and leg muscles in the same somatic segment as the innervations of the ovaries. The treatment period was 16 weeks and women received in total 14 treatments.	Improvement of: menstruation pattern, LH/FSH ratio, estrogen, and testosterone
5 E. Stener-Victorin et al., 2000.	EA is used on 24 anovulatory women with PCOS twice a week for eight times when compared with the baseline values. Moreover, acupuncture over the acupoint LI4 was reported to inhibit uterine motility.	The percentage of ovulatory cycles increased 66% up to 3 months after acupuncture

TABLE 5: EFFECTS OF RESVERATROL AND EXERCISE ON HUMANS AND RATS WITH PCOS

References	Description	Results
6 A. Benrick et al., 2013	Female Wistar rats: Resveratrol was administered 5 days a week for 4 weeks and for all 7 days of the final treatment week. Rats in the exercise group had free access to a running wheel for 5wks .	<i>Exercise and Resveratrol treatments decreased adiposity, improving estrus cyclicity in PCOS rats. Exercise is more effective than resveratrol.</i>
7 E. Stener-Victorin et al., 2009.	84 women were randomized: EA (n_33), physical exercise (n_34), or untreated control (n_17) group.	EA and physical exercise reduced high sympathetic nerve activity in PCOS.

IV. DISCUSSION

In the experiment no.1 Manni et al. demonstrated that repeated low-frequency EA treatments modulate the expression of mRNA and the amount and distribution of proteins of $\alpha 1$ -, and $\beta 2$ -adrenoreceptors (ARs), and of p75NTR in rats with steroid-induced PCOS [71].

In the study no.2 electrical and manual muscle stimulation affect glucose homeostasis through different mechanisms in rats with DHT induced PCOS. Manual stimulation improved whole-body glucose tolerance, an effect that was not observed after electrical stimulation, but did not affect molecular signaling pathways to the same extent as electrical stimulation. Although more functional signaling pathways related to insulin sensitivity were affected by electrical stimulation, maybe manual stimulation of acupuncture has a greater effect on glucose tolerance [78].

In the experiment no.3 E. Stener-Victorin et al. demonstrated high sympathetic activity in PCOS model rats and its decrease after EA treatment [73].

In the study no.4 Johansson et al. addressed acupuncture as a potential treatment option in women with PCOS. Several studies indicated in fact that acupuncture is beneficial for ovulatory dysfunction in PCOS. This is also related to decreased levels of sex steroids and inhibin B. Although the clinical data does not support changes in LH pulsatility/secretion pattern as a possible mediator for this effect, there is strong evidence of central components that probably involve both opioid and sympathetic activities, an effect that may be mediated via the androgen receptor [78].

In the experiment no.5 Stener-Victorin et al. demonstrated that the mean pulsatility index (PI) of uterine vessels was significantly reduced shortly after electroacupuncture. This may improve the endometrial environment for embryo implantation [73].

In the study no.6 A. Benrick et al. performed a 5-6 weeks resveratrol treatment on PCOS rats and did not find improvements in insulin sensitivity in DHT-induced PCOS rats, while exercise restored insulin sensitivity to a similar level as in control rats. Physical exercise also had beneficial effects on fat mass, adipocyte size, and estrus cyclicity [92].

In the experiment no. 7, it is evidenced that low-frequency EA and physical exercise decrease high sympathetic nerve activity in women with PCOS. Thus, since muscle sympathetic nerve activity (MSNA) can increase the cardiovascular risk, treatment with low-frequency EA or physical exercise aimed to reduce MSNA may be of importance for women with PCOS [98].

The effects of acupuncture on NGF expression are summarised in figure 5 [71, 73, 74, 99].

V. CONCLUSION

Several studies confirmed that acupuncture, physical activity and (to a lesser extent) resveratrol may improve metabolic, hormonal and psychological profile of PCOS in women. These positive results have been shown also in animal models, underlying the importance of these treatments, and their combinations, in clinical applications.

ACKNOWLEDGMENT

The authors thanks Dr. Luigi Manni of the Institute of Translational Pharmacology, National Research Council (CNR), Rome, Italy for the precious consultation and revision.




Fig. 5 Neurophysiological effects of acupuncture (A), EA (B) and physical activity (C) in rats affected by estradiol-induced polycystic ovaries

(A) Reprinted with modifications from Franconi et al. (2011), copyright (2011) with permission from Editrice Kurtis s.r.l.;

(B) parts of this figure were originally published in Stener-Victorin et al. (2003);

(C) part of this figure was originally published in L. Manni Ph.D. thesis (ISBN: 91-628-6484-X).

REFERENCES

- [1] C. Gilling-Smith, H. Story, V. Rogers, S. Franks. Evidence for a primary abnormality of thecal cell steroidogenesis in the polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 47: 93–99, 1997.
- [2] R. Rebar, HL. Judd, J. Rakoff, G. Vandenberg, F. Naftolin. Characterization of the inappropriate gonadotropin secretion in polycystic ovary syndrome. *J Clin Invest* 1976;57:1320–1329.
- [3] JH. Levin, E. Carmina, RA. Lobo. Is the inappropriate gonadotropin secretion of patients with polycystic ovary syndrome similar to that of patients with adult-onset congenital adrenal hyperplasia? *Fertil Steril* 1991;56:635–640.
- [4] CL. Pastor, ML. Griffin-Korf, JA. Aloï, WS. Evans, JC. Marshall. Polycystic ovary syndrome: evidence for reduced sensitivity of the gonadotropin-releasing hormone pulse generator to inhibition by estradiol and progesterone. *J Clin Endocrinol Metab* 1998;83:582–590
- [5] SK. Blank, CR. McCartney, S. Chhabra, KD. Helm, CA. Eagleson, RJ. Chang, JC. Marshall. Modulation of GnRH pulse generator sensitivity to progesterone inhibition in hyperandrogenic adolescent girls- Implications for regulation of pubertal maturation. *J Clin Endocrinol Metab* 2009;94:2360–2366.
- [6] T.D.Pache, S.Chadha, L. J. G. Gooren et al., “Ovarianmorphology in long-term androgen-treated female to male transsexuals. A human mode for the study of polycystic ovarian syndrome?” *Histopathology*, vol. 19, no. 5, pp. 445–452, 1991.
- [7] TM. Barber, MI. McCarthy, JA. Wass, S. Franks. Obesity and polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 65: 137–145, 2006.
- [8] A. Sato, Y. Sato. Regulation of regional cerebral blood flow by cholinergic fibers originating in the basal forebrain. *Neurosci Res* 14: 242–274, 1992.
- [9] S. Franks, “Do animal models of polycystic ovary syndrome help to understand its pathogenesis and management? Yes, but their limitations should be recognized,” *Endocrinology*, vol. 150, no. 9, pp. 3983–3985, 2009.
- [10] K.Hogg, J. M. Young, E. M. Oliver, C. J. Souza, A. S.McNeilly, and W. C. Duncan, “Enhanced thecal androgen production is prenatally programmed in an

- ovine model of polycystic ovary syndrome,” *Endocrinology*, vol. 153, no. 1, pp. 450–461, 2012.
- [11] T. Sir-Petermann, M. Maliqueo, B. Angel, H. E. Lara, F. P´erez-Bravo, and S. E. Recabarren, “Maternal serum androgens in pregnant women with polycystic ovarian syndrome: possible implications in prenatal androgenization,” *Human Reproduction*, vol. 17, no. 10, pp. 2573–2579, 2002.
- [12] L. Mannerås, S. Cajander, A. Holmång et al., “A new rat model exhibiting both ovarian and metabolic characteristics of polycystic ovary syndrome,” *Endocrinology*, vol. 148, no. 8, pp. 3781–3791, 2007.
- [13] E. L. van Houten, P. Kramer, A. McLuskey et al., “Reproductive and metabolic phenotype of a mouse model of PCOS,” *Endocrinology*, vol. 153, pp. 2861–2869, 2012.
- [14] A. L. L. Durlinger, P. Kramer, B. Karels et al., “Control of primordial follicle recruitment by anti-müllerian hormone in the mouse ovary,” *Endocrinology*, vol. 140, no. 12, pp. 5789–5796, 1999.
- [15] A. L. L. Durlinger, M. J. G. Gruijters, P. Kramer et al., “Anti-Müllerian hormone inhibits initiation of primordial follicle growth in the mouse ovary,” *Endocrinology*, vol. 143, no. 3, pp. 1076–1084, 2002.
- [16] H. E. Lara, J. L. Ferruz, S. Luza, D. A. Bustamante, Y. Borges, S. R. Ojeda. Activation of ovarian sympathetic nerves in polycystic ovary syndrome. *Endocrinology* 133: 2690–2695, 1993.
- [17] Y. B. Sverrisdóttir, T. Mogren, J. Kataoka, P. O. Janson, E. Stener-Victorin. Is polycystic ovary syndrome associated with high sympathetic nerve activity and size at birth? *Am J Physiol Endocrinol Metab* 294: E576–E581, 2008
- [18] S. K. Blank, C. R. McCartney, and J. C. Marshall, “The origins and sequelae of abnormal neuroendocrine function in polycystic ovary syndrome,” *Human Reproduction Update*, vol. 12, no. 4, pp. 351–361, 2006.
- [19] E. Stener-Victorin, E. Jedel and L. Mannerås (2008) Acupuncture in Polycystic Ovary Syndrome: Current Experimental and Clinical Evidence *Journal of Neuroendocrinology* 20, 290–298
- [20] D. A. Magoffin, “Ovarian theca cell,” *International Journal of Biochemistry & Cell Biology*, vol. 37, no. 7, pp. 1344–1349, 2005.
- [21] V. L. Nelson, R. S. Legro, J. F. Strauss III, and J. M. McAllister, “Augmented androgen production is a stable steroidogenic phenotype of propagated theca cells from polycystic ovaries,” *Molecular Endocrinology*, vol. 13, no. 6, pp. 946–957, 1999.
- [22] D. C. Cumming, R. L. Reid, M. E. Quigley, R. W. Rebar, S. S. Yen. Evidence for decreased endogenous dopamine and opioid inhibitory influences on LH secretion in polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 1984; 20: 643–648.
- [23] B. J. Hayden, A. H. Balen. The role of the central nervous system in the pathogenesis of polycystic ovary syndrome. *Minerva Ginecol* 2006; 58:41–54.
- [24] E. Carmina, E. C. Ditkoff, G. Malizia, A. G. Vijod, A. Janni, R. A. Lobo. Increased circulating levels of immunoreactive beta-endorphin in polycystic ovary syndrome is not caused by increased pituitary secretion. *Am J Obstet Gynecol* 1992; 167: 1819–1824.
- [25] R. A. Lobo, L. R. Granger, W. L. Paul, U. Goebelsmann, D. R. Jr. Mishell. Psychological stress and increases in urinary norepinephrine metabolites, platelet serotonin, and adrenal androgens in women with polycystic ovary syndrome. *Am J Obstet Gynecol* 1983; 145: 496–503.
- [26] X. Wu, K. Sallinen, S. Zhou, Y. Su, P. Pollanen, R. Erkkola. Androgen excess contributes to altered growth hormone/insulin-like growth factor-1 axis in nonobese women with polycystic ovary syndrome. *Fertil Steril* 2000; 73: 730–734.
- [27] J. N. Cohn, T. B. Levine, M. T. Olivari, V. Garberg, D. Lura, G. S. Francis, A. B. Simon, T. Rector. Plasma norepinephrine as a guide to prognosis in patients with chronic congestive heart failure. *N Engl J Med* 1984; 311: 819–823.
- [28] D. M. Kaye, G. W. Lambert, J. Lefkowitz, M. Morris, G. Jennings, M. D. Esler. Neurochemical evidence of cardiac sympathetic activation and increased central nervous system norepinephrine turnover in severe congestive heart failure. *J Am Coll Cardiol* 1994; 23: 570–578.
- [29] K. Patel, M. S. Coffler, M. H. Dahan, P. J. Malcom, R. Deutsch, and R. J. Chang, “Relationship of GnRH-stimulated LH release to episodic LH secretion and baseline endocrine-metabolic measures in women with polycystic ovary syndrome,” *Clinical Endocrinology*, vol. 60, no. 1, pp. 67–74, 2004.
- [30] F. J. Hayes, A. E. Taylor, K. A. Martin, and J. E. Hall, “Use of a gonadotropin-releasing hormone antagonist as a physiologic probe in polycystic ovary syndrome: assessment of neuroendocrine and androgen dynamics,” *The Journal of Clinical Endocrinology and Metabolism*, vol. 83, no. 7, pp. 2343–2349, 1998.
- [31] A. Bargiota and E. Diamanti-Kandarakis “The effects of old, new and emerging medicines on metabolic aberrations in PCOS” *Ther Adv Endocrinol Metab* (2012) 3(1) 27–47
- [32] LIU Jin-xia, DUAN Li-hui, ZHU Duan-rong, WU Jing-jing, WU Yuan-zhe. “A comparative study of cognitive function in patients with polycystic ovary syndrome before and after treatment” *Journal of Medical Postgraduates*, 2010-08
- [33] T. Takahashi, “Mechanism of acupuncture on neuromodulation in the gut—a review,” *Neuromodulation*, vol. 14, no. 1, pp. 8–12, 2011.
- [34] F. Kagitani, S. Uchida, H. Hotta, and Y. Aikawa, “Manual acupuncture needle stimulation of the rat hindlimb activates groups I, II, III and IV single afferent nerve fibers in the dorsal spinal roots,” *Japanese Journal of Physiology*, vol. 55, no. 3, pp. 149–155, 2005.
- [35] A. L. Gejervall, E. Stener-Victorin, A. Møller, P. O. Janson, C. Werner, C. Bergh. Electro-acupuncture versus conventional analgesia: a comparison of pain levels during oocyte aspiration and patients’ experiences of well-being after surgery. *Hum Reprod* 2005; 20: 728–35.

- [36] J. Sandkühler. The organization and function of endogenous antinociceptive systems. *Prog Neurobiol* 1996;1:49–81.
- [37] DJ. Mayer. Biological mechanisms of acupuncture. *Prog Brain Res* 2000;122:457–77.
- [38] WT. Zhang, Z. Jin, GH. Cui, KL. Zhang, L. Zhang, YW. Zeng, et al. Relations between brain network activation and analgesic effect induced by low vs high frequency electrical acupoint stimulation in different subjects: a functional magnetic resonance imaging study. *Brain Res* 2003;29:168–78.
- [39] M. Ferin, R. Van de Wiele. Endogenous opioid peptides and the control of the menstrual cycle. *Eur J Obstet Gynecol Reprod Biol* 1984;18: 365–73.
- [40] F. Beissner, R. Deichmann, C. Henke, and K. J. B`ar, “Acupuncture—deep pain with an autonomic dimension?” *NeuroImage*, vol. 60, no. 1, pp. 653–660, 2012.
- [41] E. Noguchi, “Acupuncture regulates gut motility and secretion via nerve reflexes,” *Autonomic Neuroscience*, vol. 156, no. 1-2, pp. 15–18, 2010.
- [42] J. Fichna, A. Janecka, J. Costentin, and J.-C. Do Rego, “The endomorphin system and its evolving neurophysiological role,” *Pharmacological Reviews*, vol. 59, no. 1, pp. 88–123, 2007.
- [43] Z.-Q. Zhao, “Neural mechanism underlying acupuncture analgesia,” *Progress in Neurobiology*, vol. 85, no. 4, pp. 355–375, 2008.
- [44] H. Akil, S. J. Watson, E. Young, M. E. Lewis, H. Khachaturian, and J. M. Walker, “Endogenous opioids: biology and function,” *Annual Review of Neuroscience*, vol. 7, pp. 223–255, 1984.
- [45] S. Andersson and T. Lundeberg, “Acupuncture— from empiricism to science functional background to acupuncture effects in pain and disease,” *Medical Hypotheses*, vol. 45, no. 3, pp. 271–281, 1995.
- [46] P. J. Jenkins and A. Grossman, “The control of the gonadotrophin releasing hormone pulse generator in relation to opioid and nutritional cues,” *Human Reproduction*, vol. 8, supplement 2, pp. 154–161, 1993.
- [47] V. G. Macefield and L. A. Henderson, “Real-time imaging of the medullary circuitry involved in the generation of spontaneous muscle sympathetic nerve activity in awake subjects,” *Human Brain Mapping*, vol. 31, no. 4, pp. 539–549, 2010.
- [48] J. F. Thayer, A. L. Hansen, E. Saus-Rose, and B. H. Johnsen, “Heart rate variability, prefrontal neural function, and cognitive performance: the neurovisceral integration perspective on self-regulation, adaptation, and health,” *Annals of Behavioral Medicine*, vol. 37, no. 2, pp. 141–153, 2009.
- [49] J.C. Hsieh, C.H. Tu, F.P. Chen, M.C. Chen, T.C. Yeh, H.C. Cheng, Y.T. Wu, R.S. Liu, L.T. Ho, 2001. Activation of the hypothalamus characterizes the acupuncture stimulation at the analgesic point in human: a positron emission tomography study. *Neurosci. Lett.* 307, 105–108.
- [50] K.K. Hui, J. Liu, A.J.W. Chen, M.T. Wu, B.R. Rosen, K.K.. Kwong, 1997. Effects of acupuncture on human limbic system and basal ganglia measured by fMRI. *NeuroImage* 5.
- [51] K.K.. Hui, J. Liu, N. Makris, R.L. Gollub, A.J. Chen, C.I. Moore, D.N. Kennedy, B.R. Rosen, K.K.. Kwong, 2000. Acupuncture modulates the limbic system and subcortical gray structures of the human brain: evidence from fMRI studies in normal subjects. *Hum. Brain Mapp.* 9, 13– 25.
- [52] J. Kong et al. 2002. A pilot study of functional magnetic resonance imaging of the brain during manual and electroacupuncture stimulation of acupuncture point (LI-4 Hegu) in normal subjects reveals differential brain activation between methods. *J. Altern. Complement. Med.* 8, 411– 419.
- [53] V. Napadow, N.Makris, , J.Liu, , N.W. Kettner, K.K. Kwong, K.K.S. Hui, 2005. Effects of electro-acupuncture vs. manual acupuncture on the human brain as measured by fMRI. *Hum. Brain Mapp.* 24 (3),193– 205
- [54] K.S. Kathleen Hui,a, et al. The integrated response of the human cerebro-cerebellar and limbic systems to acupuncture stimulation at ST 36 as evidenced by fMRI *NeuroImage* 27 (2005) 479 – 496
- [55] M. J. Smith and L. Jennes, “Neural signals that regulate GnRH neurones directly during the oestrous cycle,” *Reproduction*, vol. 122, no. 1, pp. 1–10, 2001
- [56] F. Kimura and T. Funabashi, “Two subgroups of gonadotropin-releasing hormone neurons control gonadotropin secretion in rats,” *News in Physiological Sciences*, vol. 13, no. 5, pp. 225–231, 1998.
- [57] M. Xiao-Peng, L.-Y. Tan, Y. Yang et al., “Effect of electroacupuncture on substance P, its receptor and corticotropin-releasing hormone in rats with irritable bowel syndrome,” *World Journal of Gastroenterology*, vol. 15, no. 41, pp. 5211–5217, 2009.
- [58] G. Jansen, T. Lundeberg, J. Kjartansson, and U. E. Samuelson, “Acupuncture and sensory neuropeptides increase cutaneous blood flow in rats,” *Neuroscience Letters*, vol. 97, no. 3, pp. 305–309, 1989.
- [59] M. Sandberg, T. Lundeberg, L.-G. Lindberg, and B. Gerdle, “Effects of acupuncture on skin and muscle blood flow in healthy subjects,” *European Journal of Applied Physiology*, vol. 90, no. 1-2, pp. 114–119, 2003.
- [60] M. Passatore and S. Roatta, “Influence of sympathetic nervous system on sensorimotor function: whiplash associated disorders (WAD) as a model,” *European Journal of Applied Physiology*, vol. 98, no. 5, pp. 423–449, 2006.
- [61] I. Dawidson, B. Angmar-M`ansson, M. Blom, E. Theodorsson, and T. Lundeberg, “The influence of sensory stimulation (acupuncture) on the release of neuropeptides in the saliva of healthy subjects,” *Life Sciences*, vol. 63, no. 8, pp. 659–674, 1998.
- [62] A. Sato, Y. Sato, M. Shimura, and S. Uchida, “Calcitonin gene-related peptide produces skeletal muscle vasodilation following antidromic stimulation of unmyelinated afferents in the dorsal root in rats,” *Neuroscience Letters*, vol. 283, no. 2, pp. 137–140, 2000
- [63] Z. L. Guo and J. C. Longhurst, “Expression of c-Fos in arcuate nucleus induced by electroacupuncture: relations to neurons containing opioids and glutamate,” *Brain Research*, vol. 1166, no. 1, pp. 65–76, 2007.

- [64] Z. L. Guo, A. R. Moazzami, S. Tjen-A-Looi, and J. C. Longhurst, "Responses of opioid and serotonin containing medullary raphe neurons to electroacupuncture," *Brain Research*, vol. 1229, pp. 125–136, 2008.
- [65] M. Li, S. C. Tjen-A-Looi, and J. C. Longhurst, "Electroacupuncture enhances preproenkephalin mRNA expression in rostral ventrolateral medulla of rats," *Neuroscience Letters*, vol. 477, no. 2, pp. 61–65, 2010.
- [66] H. Harbach, B. Moll, R. H. Boedeker et al., "Minimal immunoreactive plasma β -endorphin and decrease of cortisol at standard analgesia or different acupuncture techniques," *European Journal of Anaesthesiology*, vol. 24, no. 4, pp. 370–376, 2007.
- [67] S. C. Tjen-A-Looi, P. Li, and J. C. Longhurst, "Role of medullary GABA, opioids, and nociceptin in prolonged inhibition of cardiovascular sympathoexcitatory reflexes during electroacupuncture in cats," *The American Journal of Physiology*, vol. 293, no. 6, pp. H3627–H3635, 2007.
- [68] S. C. Tjen-A-Looi, P. Li, and J. C. Longhurst, "Processing cardiovascular information in the vPAG during electroacupuncture in rats: roles of endocannabinoids and GABA," *Journal of Applied Physiology*, vol. 106, no. 6, pp. 1793–1799, 2009.
- [69] L. Mannerås, I. H. Jonsdottir, A. Holmång, M. Lönn, and E. Stener-Victorin, "Low-frequency electroacupuncture and physical exercise improve metabolic disturbances and modulate gene expression in adipose tissue in rats with dihydrotestosterone-induced polycystic ovary syndrome," *Endocrinology*, vol. 149, no. 7, pp. 3559–3568, 2008.
- [70] L. Mannerås, S. Cajander, M. Lönn, and E. Stener-Victorin, "Acupuncture and exercise restore adipose tissue expression of sympathetic markers and improve ovarian morphology in rats with dihydrotestosterone-induced PCOS," *The American Journal of Physiology*, vol. 296, no. 4, pp. R1124–R1131, 2009.
- [71] L. Manni, T. Lundeberg, A. Holmång, L. Aloe, and E. Stener-Victorin, "Effect of electroacupuncture on ovarian expression of $\alpha(1)$ - and $\alpha(2)$ -adrenoceptors, and p75 neurotrophin receptors in rats with steroid-induced polycystic ovaries," *Reproductive Biology and Endocrinology*, vol. 3, article 21, 2005.
- [72] G. A. Dissen, C. Garcia-Rudaz, A. Paredes, C. Mayer, A. Mayerhofer, and S. R. Ojeda, "Excessive ovarian production of nerve growth factor facilitates development of cystic ovarian morphology in mice and is a feature of polycystic ovarian syndrome in humans," *Endocrinology*, vol. 150, no. 6, pp. 2906–2914, 2009.
- [73] E. Stener-Victorin, T. Lundeberg, U. Waldenström et al., "Effects of electroacupuncture on nerve growth factor and ovarian morphology in rats with experimentally induced polycystic ovaries," *Biology of Reproduction*, vol. 63, no. 5, pp. 1497–1503, 2000.
- [74] E. Stener-Victorin, T. Lundeberg, S. Cajander et al., "Steroid-induced polycystic ovaries in rats: effect of electroacupuncture on concentrations of endothelin-1 and nerve growth factor (NGF), and expression of NGF mRNA in the ovaries, the adrenal glands, and the central nervous system," *Reproductive Biology and Endocrinology*, vol. 1, article 33, 2003.
- [75] L. Mani, M. L. Roco, S. B. Paparo, and M. Guaragna, "Electroacupuncture and nerve growth factor: potential clinical applications," *Archives Italiennes de Biologie*, vol. 149, no. 2, pp. 247–255, 2011.
- [76] Y. Feng, J. Johansson, R. Shao, L. M. Holm, H. Billig, and E. Stener-Victorin, "Electrical and manual acupuncture stimulation affect oestrous cyclicity and neuroendocrine function in an α -dihydrotestosterone-induced rat polycystic ovary syndrome model," *Experimental Physiology*, vol. 97, no. 5, pp. 651–662, 2012.
- [77] S. Jonard and D. Dewailly, "The follicular excess in polycystic ovaries, due to intra-ovarian hyperandrogenism, may be the main culprit for the follicular arrest," *Human Reproduction Update*, vol. 10, no. 2, pp. 107–117, 2004.
- [78] J. Johansson, L. Redman, P. P. Veldhuis et al., "Acupuncture for ovulation induction in polycystic ovary syndrome: a randomized controlled trial," *American Journal of Physiology Endocrinology and Metabolism*, vol. 304, no. 9, pp. E934–E943, 2013.
- [79] G. Faulds, M. Rydén, I. Ek, H. Wahrenberg, and P. Arner, "Mechanisms behind lipolytic catecholamine resistance of subcutaneous fat cells in the polycystic ovarian syndrome," *Journal of Clinical Endocrinology and Metabolism*, vol. 88, no. 5, pp. 2269–2273, 2003.
- [80] L. Mannerås-Holm, H. Leonhardt, J. Kullberg et al., "Adipose tissue has aberrant morphology and function in PCOS: enlarged adipocytes and low serum adiponectin, but not circulating sex steroids, are strongly associated with insulin resistance," *Journal of Clinical Endocrinology and Metabolism*, vol. 96, no. 2, pp. E304–E311, 2011.
- [81] V. Skov, D. Glintborg, S. Knudsen et al., "Reduced expression of nuclear-encoded genes involved in mitochondrial oxidative metabolism in skeletal muscle of insulin-resistant women with polycystic ovary syndrome," *Diabetes*, vol. 56, no. 9, pp. 2349–2355, 2007.
- [82] J. D. Covington, E. Ravussin, and L. M. Redman, "Reduced expression of SIRT1 and SIRT3 in the skeletal muscle of PCOS women," in *The Endocrine Society's 95th Annual Meeting and Expo (ENDO '12)*, San Francisco, Calif, USA, June 2013.
- [83] K. Szkudelska and T. Szkudelski, "Resveratrol, obesity and diabetes," *European Journal of Pharmacology*, vol. 635, no. 1–3, pp. 1–8, 2010.
- [84] L. L. Chen, H. H. Zhang, J. Zheng et al., "Resveratrol attenuates high-fat diet-induced insulin resistance by influencing skeletal muscle lipid transport and subsarcolemmal mitochondrial beta-oxidation," *Metabolism*, vol. 60, pp. 1598–1609, 2011.
- [85] M. Lagogue, C. Argmann, Z. Gerhart-Hines et al., "Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1 α ," *Cell*, vol. 127, no. 6, pp. 1109–1122, 2006.
- [86] I. Ortega, J. A. Villanueva, D. H. Wong et al., "Resveratrol reduces steroidogenesis in rat ovarian theca-

interstitial cells: the role of inhibition of Akt/PKB signaling pathway," *Endocrinology*, vol. 153, pp. 4019–4029, 2012.

[87] D. H. Wong, J. A. Villanueva, A. B. Cress, and A. J. Duleba, "Effects of resveratrol on proliferation and apoptosis in rat ovarian theca-interstitial cells," *Molecular Human Reproduction*, vol. 16, no. 4, pp. 251–259, 2010.

[88] M. T. Borra, B. C. Smith, and J.M.Denu, "Mechanism of human SIRT1 activation by resveratrol," *Journal of Biological Chemistry*, vol. 280, no. 17, pp. 17187–17195, 2005.

[89] M. S. Sang, J. C. Il, and G. K. Sang, "Resveratrol protects mitochondria against oxidative stress through AMP-activated protein kinase-mediated glycogen synthase kinase-3 α inhibition downstream of poly(ADP-ribose) polymerase-LKB1 pathway," *Molecular Pharmacology*, vol. 76, no. 4, pp. 884–895, 2009.

[90] P. S. Duggal, K. H. Van Der Hoek, C. R. Milner et al., "The in vivo and in vitro effects of exogenous leptin on ovulation in the rat," *Endocrinology*, vol. 141, no. 6, pp. 1971–1976, 2000.

[91] V.W.Dolinsky, C. F. Rueda-Clausen, J. S.Morton, S.T.Davidge, and J. R. Dyck, "Continued postnatal administration of resveratrol prevents diet-induced metabolic syndrome in rat offspring born growth restricted," *Diabetes*, vol. 60, pp. 2274–2284, 2011.

[92] A. Benrick, M. Maliqueo, S.Miao, J. A. Villanueva, Yi Feng, C. Ohlsson, A. J. Duleba, and E. Stener-Victorin Resveratrol Is Not as Effective as Physical Exercise for Improving Reproductive and Metabolic Functions in Rats with Dihydrotestosterone-Induced Polycystic Ovary Syndrome. Evidence-Based Complementary and Alternative Medicine Volume 2013

[93] J. M. H. Elbers, E. J. Giltay, T. Teerlink et al., "Effects of sex steroids on components of the insulin resistance syndrome in transsexual subjects," *Clinical Endocrinology*, vol. 58, no. 5, pp. 562–571, 2003.

[94] J. Holte, T. Bergh, C. Berne, L. Wide, H. Lithell. Restored insulin sensitivity but persistently increased early insulin-secretion after weight-loss in obese women with polycystic-ovary-syndrome. *J Clin Endocrinol Metab* 1995;80:2586–2593.

[95] J-P. Despres et al., Abdominal obesity and the metabolic syndrome: contribution to global cardiometabolic risk. *Arterioscler Thromb Vasc Biol* 2008; 28:1039–1049.

[96] MM. Huber-Buchholz, DGP. Carey, RJ. Norman. Restoration of reproductive potential by lifestyle modification in obese polycystic ovary syndrome: role of insulin sensitivity and luteinizing hormone. *J Clin Endocrinol Metab* 1999;84:1470–1474.

[97] LJ. Moran, M. Noakes, PM. Clifton, L. Tomlinson, RJ. Norman. Dietary composition in restoring reproductive and metabolic physiology in overweight women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2003;88:812–819.

[98] E. Stener-Victorin, E. Jedel, Per O. Janson, and Y. Bergmann. Low-frequency electroacupuncture and physical exercise decrease high muscle sympathetic nerve activity in polycystic ovary syndrome. *Am J Physiol Regul Integr Comp Physiol* 297: R387–R395, 2009. First published June 3, 2009;oi:10.1152/ajpregu.00197.2009.

[99] Manni, L., Cajander, S., Lundeborg, T., Naylor, A. S., Aloe, L., Holmang, A., et al. (2005). Effect of exercise on ovarian morphology and expression of nerve growth factor and alpha(1)- and beta(2)-adrenergic receptors in rats with steroid-induced polycystic ovaries. *Journal of Neuroendocrinology*, 17(12), 846–858.