

Università degli Studi di Salerno
DIPARTIMENTO DI SCIENZE ECONOMICHE E STATISTICHE

Pietro Coretto ¹ Christian Hennig ²

IDENTIFIABILITY FOR MIXTURES OF
DISTRIBUTIONS FROM A LOCATION-SCALE
FAMILY WITH UNIFORMS

WORKING PAPER 3.186

¹Dipartimento di Scienze Economiche e Statistiche Università di Salerno, Italy
pcoretto@unisa.it

²Department of Statistical Sciences University College London, UK
chrish@stats.ucl.ac.uk

Contents

1	<i>Introduction</i>	7
2	<i>Identifiability of mixture distributions</i>	9
3	<i>Model definitions</i>	11
4	<i>Identifiability of "heterogeneous" mixtures</i>	12
5	<i>The identifiability of the model with uniform noise</i>	15
6	<i>Conclusions</i>	19

Abstract

In this paper we study the indentifiability of a class of mixture models where a finite number of one-dimensional location scale distributions is mixed with a finite number of uniform distributions on an interval. We define identifiability and we show that, under certain conditions, the afore-mentioned class of distributions is identifiable.

Keywords

Mixture models, identifiability, robustness

1 Introduction

Mixture models are widely used in many contexts. In cluster analysis and classification, mixture distributions have the role to model data with group-structures. Banfield and Raftery (1993) called this approach “model based clustering”. Several different mixture models have been studied and estimation methods proposed over the years. Normal mixtures and the related theory of maximum likelihood estimation played an important role in the research. However, maximum likelihood estimates for a wide class of location-scale mixtures are not robust (see Hennig, 2004). Fraley and Raftery (1998) treats outliers as points coming from some unknown mixture component interpreted as “noise”. They proposed a model where a uniform distribution is mixed with the Gaussians in order to account for noise. Hennig (2004) defined robustness measures for clusters, and he studied the properties of several robust model based clustering methods including Gaussian mixtures with uniform noise, uniform-normal mixture from now onward. However, identifiability for this model has not been studied. Teicher (1963) shows that univariate normal mixtures are identifiable, while in general mixture of uniform distributions are not. The proposal of Banfield and Raftery (1993) consists of a mixture with arbitrary number of normals with the addition of a uniform component, the authors proposed estimation methods as well as a computational procedures to get estimates.

The aim of this paper is to define and show identifiability for a general class of models which includes the afore-mentioned uniform-normal mixture. In a nutshell, in the context of parametric family of distributions, identifiability means that different parameter values lead to different distributions. If a dis-

tribution is not identifiable, the same parameter can define more than one distribution; this would affect any consistency statement about estimators. Identifiability is also relevant from the practical point of view. In fact, if a population is represented by some non-identifiable parametric distribution this means that the population under study can be represented by more than one parameter. Here we do not constraint the number of uniform components, and we do not confine ourself to the case of Gaussian mixtures. Our model of interest will be a finite mixture of location-scale distributions mixed with a finite number of uniform distributions. This means that we have a mixture with components belonging to different families. The original definition of identifiability for mixtures was given by Teicher (1961). This definition does not take into account the situation where distributions from different families are mixed and it is required that the family memberships of the components are identified. For the class of models under study we want to identify the parameters, the number of components and their family memberships. Thus we will propose a definition of identifiability which extends the one proposed by Teicher (1961). Yakowitz and Spragins (1968) and Atienza, Garcia-Heras, and Munoz-Pichardo (2006) provided sufficient conditions to show identifiability under Teicher's definition. However, The afore-mentioned sufficient conditions are not applicable in our situation. Heuristically, the idea under our identifiability proof is simple. The distribution function of the class of mixture models under study is not differentiable at a set of points which coincides with the set of the uniform parameters. Thus the points where the mixture distribution is not differentiable will identify the uniform parameters; then we remove the uniform components and we will identify the remaining components. Even though the idea behind this identifiability results is simple, the formal implementation is not easy.

The paper is organized as follows: first we introduce motivations for the class of models under study; in section 2 we introduce the general definition of identifiability given by Teicher (1961); in section 3 we define the class of models under study and in section 4 we extend the Teicher's definition of identifiability; in section 5 we state and prove some identifiability results for the class of models under study; finally in section 6 we draw conclusions and outline some future works.

2 Identifiability of mixture distributions

Let \mathcal{P}_θ be a family of probability measures indexed by some unknown – finite or infinite dimensional – index $\theta \in \Theta$ which we call parameter. We observe an experiment generated by some member of \mathcal{P}_θ . The main problem of statistical inference is to infer θ based on observed data. Instead, identification is a pre-inferential problem which is devoted to assess whether with data at hand it is possible to state that different parameter values correspond to different probability measures $P \in \mathcal{P}_\theta$, where the meaning of the word “different” has to be specified. Roughly speaking identifiability means that there exists a sort of one-to-one correspondence between the indexes $\theta \in \Theta$ and $P \in \mathcal{P}_\theta$. The first account of identification of mixture models was given by Feller (1943) and since then many results extended that work in several directions (we shall review those results in the following paragraphs).

Identifiability is a general concept that has to be carefully defined depending on the context. The very first definition of identifiability for finite mixtures was formalized first by Teicher (1961). Let $\mathcal{F} := \{F(x; \theta) : x \in \mathbb{R}^q, \theta \in \mathbb{R}^k\}$ be a family of distribution functions over \mathbb{R}^q indexed by a point θ in a Borel subset of \mathbb{R}^k such that $F(\cdot; \theta)$ is measurable on $\mathbb{R}^q \times \mathbb{R}^k$. Let $G \in \mathcal{G}$ be a k -dimensional distribution function with the underlying measure assigning total mass to \mathbb{R}^k . Let \mathcal{H} be a family of distribution functions. We consider a map $Q : \mathcal{G} \rightarrow \mathcal{H}$, where its image is defined as $Q(G) = H$, $H(x) = \int_{\mathbb{R}^k} F(x; \theta) dG(\theta)$. Following Teicher (1961), the mixture model generated by the family \mathcal{F} with mixing distribution in \mathcal{G} is said to be identifiable if given $F \in \mathcal{F}$, then Q is a one-to-one map of \mathcal{G} onto \mathcal{H} . As we have already noticed, when G is discrete, the set of all finite mixtures \mathcal{H} of the family \mathcal{F} is simply the convex hull of \mathcal{H} . Identifiability of the mixture models means that the convex hull \mathcal{F} has a uniqueness representation property which can be translated into following:

Definition 1 (Single Family Identifiability). Let \mathcal{H} be the class of finite mix-

tures generated by the class \mathcal{F} with discrete mixing distribution. Given

$$H(x, \eta) = \sum_{j=1}^s \pi_j F(x; \theta_j), \quad \pi_j > 0, \theta_j \neq \theta_r \quad \forall j, r = 1, 2, \dots, s, \quad j \neq r,$$

and

$$H(x, \eta^*) = \sum_{i=1}^t \pi_i^* F(x; \theta_i^*), \quad \pi_i^* > 0, \theta_i^* \neq \theta_k^* \quad \forall i, k = 1, 2, \dots, t, \quad i \neq k;$$

if $H(\cdot, \eta) = H(\cdot, \eta^*)$ implies that $s = t$, and there is some permutation \bar{j} of the indexes $j = 1, 2, \dots, s$ such that $\pi_j = \pi_{\bar{j}}^*$ and $\theta_j = \theta_{\bar{j}}^*$, then we say that \mathcal{F} generates identifiable finite mixture distributions.

The definition above has been used to study the identification of a number of models. The wording “single family identifiability” will be clearer thereafter. Feller (1943) started the literature about identifiability of mixtures studying gamma mixture models. Teicher (1961) formalized the definition of identifiability for general mixture models. He extended the results in Feller (1943) showing the identifiability of finite mixtures generated by Poisson distributions. He also showed that models based on mixtures of uniform and binomial distributions are not identifiable. Teicher (1963) gave a sufficient condition for identifiability of a general class of finite mixture models and showed that mixtures based on univariate Gaussian distributions are identifiable. Yakowitz and Spragins (1968) defined identifiability for classes of finite mixtures (Definition 1) and gave a necessary and sufficient condition for the identifiability of such models. The main theorem in Yakowitz and Spragins (1968) states that given a discrete mixing distribution the class \mathcal{F} generates identifiable mixtures if and only if \mathcal{F} is a linearly independent set over the field of the real numbers. They apply their theory showing that exponential distributions, multivariate Gaussian distributions, Cauchy distributions and negative binomials generate identifiable mixture models. Atienza, Garcia-Heras, and Munoz-Pichardo (2006) weakened the assumptions of the sufficient conditions given by Teicher (1963) and showed that mixtures of Log-Gamma distributions and mixtures of Lognormal, Gamma and Weibull distributions are identifiable with the respect to the Definition 1.

3 Model definitions

In this section we introduce the notation and the main assumptions about the general model under study. Let $0 < s < \infty$ be the number of components in our mixture distribution, and let q be the number of uniform components $0 < q < s$ in the mixture. Let X be a real valued random variable distributed according to the following distribution function:

$$G(x; \eta) = \sum_{k=1}^q \pi_k U(x; \theta_k) + \sum_{l=q+1}^s \pi_l \Phi(x; \theta_l), \quad (3.1)$$

where $\eta = (\pi, \theta)$, $\pi = (\pi_1, \pi_2, \dots, \pi_s)$, $0 < \pi_j < 1$, $\sum_{j=1}^s \pi_j = 1$. Here $\theta = (\theta_1, \theta_2, \dots, \theta_s)$, where $\theta_k = (a_k, b_k)$, a_k and b_k take values on the real line, and $-\infty < a_k < b_k < +\infty$ for each $k = 1, 2, \dots, q$. Thus $\pi \in (0, 1)^s$, $\theta_k \in \Theta_1 := \mathbb{R}^{2q}$ for $k = 1, 2, \dots, q$. The parameter θ_l lies in some finite dimensional space Θ_2 for each $l = q+1, q+2, \dots, s$. Furthermore the parameter space is denoted by $\Gamma := (0, 1)^s \times \mathbb{R}^{2q} \times \Theta_2^{s-q}$. U is the uniform distribution function, i.e.

$$U(x; \theta_k) = \frac{x - a_k}{b_k - a_k} \mathbf{1}_{[a_k, b_k]}(x) + \mathbf{1}_{(b_k, +\infty)}(x),$$

$k = 1, 2, \dots, q$, with $\mathbf{1}_A$ being the indicator function of the set A . The distribution function U has the density

$$u(x; \theta_k) = \frac{\mathbf{1}_{[a_k, b_k]}(x)}{b_k - a_k}.$$

The distribution function Φ belongs to a family of distributions satisfying

Assumption 1. $\Phi(x; \theta)$, $\theta \in \Theta_2$, is absolutely continuous with the respect to Lebesgue measure. It has density $\phi(x; \theta)$, $\theta \in \Theta_2$, which is continuous both with the respect to $x \in \mathbb{R}$ and $\theta \in \Theta_2$.

For notational convenience we will often rewrite the model in (3.1) as

$$G(x, \eta) := \sum_{j=1}^s \pi_j F_{v_j}(x; \theta_j) \quad (3.2)$$

where $v_j = \{1, 2\}$ for $j = 1, 2, \dots, s$, when $v_j = 1$ then $F_{v_j} = U$, whenever $v_j = 2$ then $F_{v_j} = \Phi$. Moreover $g(x; \eta)$ will denote the density of $G(x; \eta)$.

4 Identifiability of “heterogeneous” mixtures

In section 2 we introduced the identifiability problem for general finite mixtures. In this section we define and study the identifiability of a class of models which consists of a mixture of distributions coming from different families. We are in a situation where a finite number of distributions belonging to a general class of continuous distributions is mixed with a finite number of uniform distributions. We call such a mixture distribution: heterogeneous. Here the wording heterogeneous mixtures means that the components in the mixture belong to different families of distributions. Such a statistical model can be very attractive in all those situations where the underlying heterogeneity in the data generating process is strong enough to let us consider that groups of observations come from populations with completely different features. In fact the uniform distribution here is introduced as a probabilistic model for noise, while Φ should represent the probabilistic structure of the population under study. Here we do not require that the number of components is known, nor do we require that the number of components belonging to each of the families of distributions is known. This situation is somehow more general than that of the model proposed by Fraley and Raftery (1998), where the number of uniform components is considered as fixed and known. In fact in the uniform–normal mixture model proposed by Fraley and Raftery (1998) the number of uniform components is fixed to be one.

We now refer to section 2 where we presented the definition of identifiability as given by Teicher (1961). Let us assume that \mathcal{F}_k , with $k = 1, 2, \dots, m$, are all families of probability distribution functions. For each k , $F_k(x; \theta) \in \mathcal{F}_k$, $\theta \in \Theta_k$, and Θ_k is some finite dimensional parameter space. A general element of the set of finite mixtures generated by the class $\mathcal{E} = \cup_{k=1}^m \mathcal{F}_k$ will be called heterogenous mixture distribution. Teicher’s definition of identifiability does not require that the number of components in

each family is identified. However this is relevant in a situation where membership to different population components have different meaning. In our model for example we want to distinguish between noise components and non-noise components, and we want that the number of distributions belonging to each of the family composing the mixture is identified. To see why definition 1 does not take into account the identifiability of family memberships let us consider some results in the paper by Atienza, Garcia-Heras, and Munoz-Pichardo (2006). The authors studied the identifiability a model proposed by Marrazzi, Paccaud, Ruffieux, and Beguin (1998) in the context of fitting the length of stay in a hospital; the model is a mixture of three components: one Lognormal, one Gamma and one Weibull distribution. Atienza, Garcia-Heras, and Munoz-Pichardo (2006) gave a new sufficient condition for identifiability of finite mixtures following Teicher's definition, and based on this they showed the identifiability of the afore-mentioned class of mixtures. However, following the proof of their theorem 3 it is clear that for some value of the parameter space, a component having Gamma distribution cannot be distinguished from a component having Weibull distribution, thus the number of components belonging to each family cannot be identified.

Here we will give a definition of identifiability which is similar to the one given by Teicher (1961) but it adds some more restrictions so that the family membership of components is taken into account in the sense explained above. It should be now clear why we named the identifiability defined by Teicher as "single family identifiability". Before we give our definition, let us introduce some more notations.

We will consider the set of all heterogenous finite mixtures generated by \mathcal{E} with discrete mixing distribution. Let $s < +\infty$ be the number of components of the heterogenous mixture, and let $c = \{n_1, n_2, \dots, n_m\}$ be a set of natural numbers where $n_k, k = 1, 2, \dots, m$, indicates the number of distributions belonging to \mathcal{F}_k being present in the mixture. From now onward it is understood that c is finite, and of course it must be $s = \sum_{k=1}^m n_k$. We will denote c as the "composition" index. \mathcal{H} is the family of all the finite mixtures generated from \mathcal{E} with discrete mixing distribution. A general element of \mathcal{H} will be $H_c(x; \eta) = \sum_{j=1}^s \pi_j F_{k_j}(x; \theta_j)$, where $k_j \in \{1, 2, \dots, m\}$

for $j = 1, 2, \dots, s$, is the index which expresses the “family membership” of the j th component (e.g. $k_2 = 1$ means that the distribution of the second mixture component belongs to \mathcal{F}_1). The parameter η lies in the parameter set Ω , and $\eta = (\pi_1, \dots, \pi_s, \theta_1, \dots, \theta_s)$. We will consider the following definition:

Definition 2 (Global Identifiability). Let \mathcal{H} the class of finite mixtures generated by the class \mathcal{E} . Let $\mathcal{H}^* \subseteq \mathcal{H}$, and $H_c \in \mathcal{H}^*$. Given

$$H_c(x, \eta) = \sum_{j=1}^s \pi_j F_{v_j}(x; \theta_j), \pi_j > 0, \theta_j \neq \theta_r \quad \forall j, r = 1, 2, \dots, s, \quad j \neq r,$$

and

$$H_{c^*}(x, \eta^*) = \sum_{j=1}^z \pi_j^* F_{v_j}(x; \theta_j^*), \pi_j^* > 0, \theta_j^* \neq \theta_k^* \quad \forall j, k = 1, 2, \dots, z, \quad j \neq k;$$

if $H_c(\cdot, \eta) = H_{c^*}(\cdot, \eta^*)$ implies $s = z$, and that there exists a permutation \bar{j} of the indexes $j = 1, 2, \dots, s$ such that $\pi_j = \pi_{\bar{j}}^*$, $\theta_j = \theta_{\bar{j}}^*$, $k_j = k_{\bar{j}}$, for $k_j, k_{\bar{j}} \in \{1, 2, \dots, m\}$, and $c = c^*$, then we say that \mathcal{E} generates globally identifiable finite mixture distributions in \mathcal{H}^* .

As highlighted before, we use the wording *global identifiability* to make a distinction between the notion of identifiability given in definition 2 with the one given in definition 1 which refers to Teicher’s definition. With reference to definition 2, we require that the permutation of the component label (the index j) is constructed so that for each family \mathcal{F}_k we identify the parameters, obtaining $\pi_j F_{k_j}(x; \theta_j) = \pi_{\bar{j}}^* F_{k_{\bar{j}}}(x; \theta_{\bar{j}}^*)$, and at the same time we require that the number of distributions identified in the family \mathcal{F}_k is consistent with the composition index c . To see the relevance of this argument let us refer to the the model proposed by Banfield and Raftery (1993). In that case we require that not only the uniform parameters, the Gaussian parameters and all proportions are identified but we also require that it is also possible to identify the number of noise components and Gaussian components.

5 The identifiability of the model with uniform noise

First, we will introduce some notations and assumptions and we also reconcile the exposition here with the notation used in the previous sections. We consider the model defined in section 3 with the addition of the following definitions: (i) \mathcal{F}_1 is the family of all uniform distributions with support on an interval; (ii) \mathcal{F}_2 is a family of one dimensional distributions satisfying assumption 1 and finite mixtures generated by \mathcal{F}_2 are identifiable in the sense of the definition 2.

Let $n_1 = q, n_2 = s - q$ and let $c = \{q, s - q\}$ be the composition index. \mathcal{H} is the family of finite mixtures generated by $\mathcal{E} = \mathcal{F}_1 \cup \mathcal{F}_2$, obtained by mixing q distributions from \mathcal{F}_1 and $s - q$ distributions from \mathcal{F}_2 . The function $g_c(x; \eta)$ will denote the density of $G_c(x; \eta)$ likewise the model defined in section 3. $G_c(x; \eta)$ is an element of \mathcal{H} , with $\eta \in \Gamma$. $\mathcal{H}^* \subset \mathcal{H}$ is the set of mixtures generated by \mathcal{E} such that if $G_c(x; \eta)$ belongs to \mathcal{H}^* , then $[a_t, b_t] \cap [a_r, b_r] = \emptyset$ for all $r, t = 1, 2, \dots, q$ and $r \neq t$.

To show identifiability here we will make use of arguments based on derivatives so that it is necessary to introduce some more notations before to state and prove the next result. We notice that the density $g_c(x; \eta)$ it is discontinuous at a finite number of points, namely at $x \in W := \{a_1, b_1, a_2, b_2, \dots, a_q, b_q\}$. Thus by properties of the Riemann integral, $dG_c(x; \eta)/dx = g_c(x; \eta)$ at all $x \in \mathbb{R} \setminus W$. However, right and left derivatives of G_c at all points in W exist and can be found by taking right and left limits of derivative quotients. The notation $D_y^-(\eta)$ and $D_y^+(\eta')$ stands for the left and right derivative of G_c respectively, and these derivatives are evaluated at a point y when the parameter vector is η' , i.e.

$$D_y^-(\eta') = \lim_{t \uparrow 0} \frac{G_c(y + t; \eta') - G_c(y; \eta')}{t},$$

$$D_y^+(\eta') = \lim_{t \downarrow 0} \frac{G_c(y + t; \eta') - G_c(y; \eta')}{t}.$$

Computing these derivatives for the model (3) and for $h = 1, 2, \dots, q$ will

give us

$$D_{a_h}^-(\eta) = \sum_{l=q+1}^{s-q} \pi_l \phi(a_h; \theta_l);$$

$$D_{a_h}^+(\eta) = \frac{\pi_h}{b_h - a_h} + \sum_{l=q+1}^{s-q} \pi_l \phi(a_h; \theta_l);$$

$$D_{b_h}^-(\eta) = \frac{\pi_h}{b_h - a_h} + \sum_{l=q+1}^{s-q} \pi_l \phi(b_h; \theta_l);$$

$$D_{b_h}^+(\eta) = \sum_{l=q+1}^{s-q} \pi_l \phi(b_h; \theta_l).$$

Theorem 1. *The class $\mathcal{E} = \mathcal{F}_1 \cup \mathcal{F}_2$ generates globally identifiable heterogeneous mixtures in $\mathcal{H}^* \subset \mathcal{H}$.*

Proof. Let us assume that $G_c(x; \eta) = G_{c^*}(x; \eta^*)$, i.e.

$$\sum_{j=1}^s \pi_j F_{v_j}(x; \theta_j) = \sum_{j=1}^z \pi_j^* F_{v_j}(x; \theta_j^*), \quad (5.1)$$

for every $x, v_j \in \{1, 2\}$ and $j = 1, 2, \dots, s, \dots, z$, i.e. without loss of generality we assume that $s \leq z$. For a given function $f(y, z)$ differentiable at least on a subset of its own domain, we define the set

$$S_f(z) := \left\{ y : \frac{\partial^-}{\partial y} f(y, z) \neq \frac{\partial^+}{\partial y} f(y, z) \right\};$$

provided that all at points in $S_f(z)$ left and right partial derivatives of f exist. The assumption that $G_c(x; \eta) = G_{c^*}(x; \eta^*)$ implies that $S_{G_c}(\eta) =$

$S_{G_{c^*}}(\eta^*)$. If $\#(A)$ stands for the cardinality of the set A , then $\#(S_{G_c}(\eta)) = \#(S_{G_{c^*}}(\eta^*)) = 2q$ which means that the number of the uniform components q is uniquely identified. Given a finite set $A := \{y_1, y_2, \dots, y_n\}$, with $y_i \in \mathbb{R}$ all $i = 1, 2, \dots, n$, $\mu(A) \in \mathbb{R}^n$ denotes a vector where the components are all the elements of A , furthermore $\bar{\mu}(A)$ is defined as $\bar{\mu}(A) = (y_{(1)}, y_{(2)}, \dots, y_{(n)})$ where $y_{(i)}$ is such that $y_{(i)} \leq y_{(i+1)}$ all $i = 1, 2, \dots, n - 1$. Now, $\bar{\mu}(S_{G_c}(\eta)) = (x_{(1)}, x_{(2)}, \dots, x_{(2q)}) = \bar{\mu}(S_{G_{c^*}}(\eta^*)) = (x_{(1)}^*, x_{(2)}^*, \dots, x_{(2q)}^*)$. We take a set of pairwise different indexes $r_i \in \{1, 2, \dots, s\}$ with $i = 1, 2, \dots, q$ and we fix

$$\begin{aligned}\theta_{r_1} &= (a_{r_1}, b_{r_1}) = (x_{(1)}, x_{(2)}), \\ \theta_{r_2} &= (a_{r_2}, b_{r_2}) = (x_{(3)}, x_{(4)}), \\ &\vdots \\ \theta_{r_q} &= (a_{r_q}, b_{r_q}) = (x_{(2q-1)}, x_{(2q)}).\end{aligned}$$

Let us take another set of pairwise different indexes $t_i \in \{1, 2, \dots, s\}$ with $i = 1, 2, \dots, q$ and we fix

$$\begin{aligned}\theta_{t_1}^* &= (a_{t_1}^*, b_{t_1}^*) = (x_{(1)}^*, x_{(2)}^*), \\ \theta_{t_2}^* &= (a_{t_2}^*, b_{t_2}^*) = (x_{(3)}^*, x_{(4)}^*), \\ &\vdots \\ \theta_{t_q}^* &= (a_{t_q}^*, b_{t_q}^*) = (x_{(2q-1)}^*, x_{(2q)}^*).\end{aligned}$$

It is clear that $\bar{\mu}(S_{G_c}(\eta)) = \bar{\mu}(S_{G_{c^*}}(\eta^*))$ implies that $\theta_{r_i}^* = \theta_{t_i}^*$ all $i = 1, 2, \dots, q$. Let us consider the equation

$$\left(D_{a_{r_i}}^+(\eta) - D_{a_{r_i}}^-(\eta)\right) (b_{r_i} - a_{r_i}) = \left(D_{a_{t_i}^*}^+(\eta^*) - D_{a_{t_i}^*}^-(\eta^*)\right) (b_{t_i}^* - a_{t_i}^*),$$

for all $i = 1, 2, \dots, q$. These equations imply that $\pi_{r_i} = \pi_{t_i}^*$ all $i = 1, 2, \dots, q$. Hence, we have that there exists a permutation \bar{j} of the indexes $j = 1, 2, \dots, s, \dots, z$ such that if $j = r_i$ then $\bar{j} = t_i$, for which

$\theta_j = (a_j, b_j) = \theta_j^* = (a_j^*, b_j^*)$, $\pi_j = \pi_j^*$ and $v_j = v_j^* = 1$. By this we have identified the number of uniform components, and all their parameters. Without loss of generality let us assume that $r_i = t_i$ for all $i = 1, 2, \dots, q$ and that $r_1, r_2, \dots, r_q = 1, 2, \dots, q$.

For $j = q + 1, q + 2, \dots, s, \dots, z$ all the mixture components belong to \mathcal{F}_2 and q is identified as well. We consider the one-to-one transformation $\tilde{\pi}_j = \pi_j / (1 - \sum_{j=1}^q \pi_j)$ for $j = q + 1, q + 2, \dots, s, \dots, z$; and $\tilde{\pi}_j^*$ is defined analogously. Note that the denominator of $\tilde{\pi}_j$ is identified, in fact it depends on $\pi_1, \pi_2, \dots, \pi_q$ which have been already identified. By (5.1) and the previous results we can write

$$\sum_{j=q+1}^s \tilde{\pi}_j F_{v_j}(x, \theta_j) = \sum_{j=q+1}^z \tilde{\pi}_j^* F_{v_j^*}(x, \theta_j^*). \quad (5.2)$$

By assumption the class of finite mixtures over \mathcal{F}_2 is identifiable with respect to definition 1, thus we have that: (i) $s = z$ and these indexes are identified; (ii) there exists some permutation \bar{j} of indexes $j = q + 1, q + 2, \dots, s$ such that $\tilde{\pi}_j = \tilde{\pi}_{\bar{j}}^*$ and $\theta_j = \theta_{\bar{j}}^*$. But, $\tilde{\pi}_j = \tilde{\pi}_{\bar{j}}^*$ implies $\pi_j = \pi_{\bar{j}}^*$. Thus the $s - q$ components belonging to \mathcal{F} , their parameters and their mixing proportions are identified. The proof is completed noting that having identified q and s it also results that $c = c^*$. \square

Given the proposition above we can easily get the next result.

Corollary 1. *Let \mathcal{F}_2 be the class of Gaussian distributions, then the class $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2$ generates globally identifiable mixtures in \mathcal{H} .*

Proof. The result follows easily by noting that: (i) Gaussian distributions clearly satisfy assumption 1; (ii) they are partially identifiable by theorem 3 in Yakowitz and Spragins (1968). \square

We defined \mathcal{F}_1 so that it contains uniform distributions having not intersecting support. The reason for this should be clear. Let us assume that \mathcal{F}_1 contains all uniform distribution with support on a real interval, and let us consider the following mixture distribution

$$\frac{1}{3}U(x; 0, 2) + \frac{1}{3}U(x; 2, 4) + \frac{1}{3}F(x; \theta),$$

where F is some distribution function satisfying assumption 1. We notice that $1/3U(x; 0, 2) + 1/3U(x; 2, 4) = 2/3U(x; 0, 4)$ so that identifiability does not hold. In fact, not only the parameters of the uniform distributions are not identifiable but also the composition index referring to the uniform components would not be identifiable.

6 Conclusions

In this paper we studied the identifiability for some class of mixture distributions. We gave a new definition of identifiability which take into account heterogeneity in the mixture, and we showed that a wide class of mixture with uniform components are identifiable. In the literature the model consisting of a Gaussian-uniform mixture have been proposed to overcome problems of robustness (see Fraley and Raftery, 1998). In this proposal the number of uniform components is fixed to be one, but we could be interested in determining whether the number of noise components is one or zero. The estimation of the number of noise components requires that it is identified. We extended this model to the case when more than one uniform components is added to a mixture of a class of continuous location-scale mixtures and we showed that the resulting mixture is identifiable. In a future paper we will study the estimation of such a model and we will explore the related computational issues.

References

- ATIENZA, N., J. GARCIA-HERAS, AND J. M. MUNOZ-PICHARDO (2006): "A new condition for the identifiability of finite mixture distributions," *Metrika*, 63, 215–221.
- BANFIELD, J., AND A. E. RAFTERY (1993): "Model-based Gaussian and non-Gaussian clustering," *Biometrics*, 49, 803–821.
- FELLER, W. (1943): "On a general class of "contagious" distributions," *The Annals of Mathematical Statistics*, 14, 389–399.
- FRALEY, C., AND A. E. RAFTERY (1998): "How many clusters? Which clus-

- tering method? Answers via model-based cluster analysis," *The Computer Journal*, 41, 578–588.
- HENNIG, C. (2004): "Breakdown Points For Maximum Likelihood Estimators of LocationScale Mixtures," *The Annals of Statistics*, 32(4), 1313–1340.
- MARRAZZI, A., F. PACCAUD, C. RUFFIEUX, AND C. BEGUIN (1998): "Fitting the distribution of lenght of stay by paramteric models," *Medical–Care*, 36(6), 915–927.
- TEICHER, H. (1961): "Identifiability of mixtures," *The Annals of Mathematical Statistics*, 32, 244–248.
- (1963): "Identifiability of finite mixtures," *The Annals of Mathematical Statistics*, 34, 1265–1269.
- YAKOWITZ, S. J., AND J. SPRAGINS (1968): "On the identifiability of finite mixtures," *The Annals of Mathematical Statistics*, 39, 209–214.

WORKING PAPERS DEL DIPARTIMENTO

- 1988, 3.1 Guido CELLA
Linkages e moltiplicatori input-output.
- 1989, 3.2 Marco MUSELLA
La moneta nei modelli di inflazione da conflitto.
- 1989, 3.3 Floro E. CAROLEO
Le cause economiche nei differenziali regionali del tasso di disoccupazione.
- 1989, 3.4 Luigi ACCARINO
Attualità delle illusioni finanziarie nella moderna società.
- 1989, 3.5 Sergio CESARATTO
La misurazione delle risorse e dei risultati delle attività innovative: una valutazione dei risultati dell'indagine CNR- ISTAT sull'innovazione tecnologica.
- 1990, 3.6 Luigi ESPOSITO - Pasquale PERSICO
Sviluppo tecnologico ed occupazionale: il caso Italia negli anni '80.
- 1990, 3.7 Guido CELLA
Matrici di contabilità sociale ed analisi ambientale.
- 1990, 3.8 Guido CELLA
Linkages e input-output: una nota su alcune recenti critiche.
- 1990, 3.9 Concetto Paolo VINCI
I modelli econometrici sul mercato del lavoro in Italia.
- 1990, 3.10 Concetto Paolo VINCI
Il dibattito sul tasso di partecipazione in Italia: una rivisitazione a 20 anni di distanza.
- 1990, 3.11 Giuseppina AUTIERO
Limiti della coerenza interna ai modelli con la R.E.H..
- 1990, 3.12 Gaetano Fausto ESPOSITO
Evoluzione nei distretti industriali e domanda di istituzione.
- 1990, 3.13 Guido CELLA
Measuring spatial linkages: input-output and shadow prices.
- 1990, 3.14 Emanuele SALSANO
Seminari di economia.

- 1990, 3.15 Emanuele SALSANO
Investimenti, valore aggiunto e occupazione in Italia in contesto biregionale: una prima analisi dei dati 1970/1982.
- 1990, 3.16 Alessandro PETRETTO- Giuseppe PISAURO
Uniformità vs selettività nella teoria della ottima tassazione e dei sistemi tributari ottimali.
- 1990, 3.17 Adalgiso AMENDOLA
Inflazione, disoccupazione e aspettative. Aspetti teorici dell'introduzione di aspettative endogene nel dibattito sulla curva di Phillips.
- 1990, 3.18 Pasquale PERSICO
Il Mezzogiorno e le politiche di sviluppo industriale.
- 1990, 3.19 Pasquale PERSICO
Priorità delle politiche strutturali e strategie di intervento.
- 1990, 3.20 Adriana BARONE - Concetto Paolo VINCI
La produttività nella curva di Phillips.
- 1990, 3.21 Emiddio GALLO
Varianze ed invarianze socio-spaziali nella transizione demografica dell'Italia post-industriale.
- 1991, 3.22 Alfonso GAMBARDELLA
I gruppi etnici in Nicaragua. Autonomia politica ed economica.
- 1991, 3.23 Maria SCATTAGLIA
La stima empirica dell'offerta di lavoro in Italia: una rassegna.
- 1991, 3.24 Giuseppe CELI
La teoria delle aree valutarie: una rassegna.
- 1991, 3.25 Paola ADINOLFI
Relazioni industriali e gestione delle risorse umane nelle imprese italiane.
- 1991, 3.26 Antonio e Bruno PELOSI
Sviluppo locale ed occupazione giovanile: nuovi bisogni formativi.
- 1991, 3.27 Giuseppe MARIGLIANO
La formazione del prezzo nel settore dell'intermediazione commerciale.
- 1991, 3.28 Maria PROTO
Risorse naturali, merci e ambiente: il caso dello zolfo.
- 1991, 3.29 Salvatore GIORDANO
Ricerca sullo stato dei servizi nelle industrie del salernitano.

- 1992, 3.30 Antonio LOPES
Crisi debitoria e politiche macroeconomiche nei paesi in via di sviluppo negli anni 80.
- 1992, 3.31 Antonio VASSILLO
Circuiti economici semplici, complessi, ed integrati.
- 1992, 3.32 Gaetano Fausto ESPOSITO
Imprese ed istituzioni nel Mezzogiorno: spunti analitici e modalità di relazione.
- 1992, 3.33 Paolo COCCORESE
Un modello per l'analisi del sistema pensionistico.
- 1994, 3.34 Aurelio IORI
Il comparto dei succhi di agrumi: un caso di analisi interorganizzativa.
- 1994, 3.35 Nicola POSTIGLIONE
Analisi multicriterio e scelte pubbliche.
- 1994, 3.36 Adriana BARONE
Cooperazione nel dilemma del prigioniero ripetuto e disoccupazione involontaria.
- 1994, 3.37 Adriana BARONE
Le istituzioni come regolarità di comportamento.
- 1994, 3.38 Maria Giuseppina LUCIA
Lo sfruttamento degli idrocarburi offshore tra sviluppo economico e tutela dell'ambiente.
- 1994, 3.39 Giuseppina AUTIERO
Un'analisi di alcuni dei limiti strutturali alle politiche di stabilizzazione nei LCDs.
- 1994, 3.40 Bruna BRUNO
Modelli di contrattazione salariale e ruolo del sindacato.
- 1994, 3.41 Giuseppe CELI
Cambi reali e commercio estero: una riflessione sulle recenti interpretazioni teoriche.
- 1995, 3.42 Alessandra AMENDOLA, M. Simona ANDREANO
The TAR models: an application on italian financial time series.
- 1995, 3.43 Leopoldo VARRIALE
Ambiente e turismo: Parco dell'Iguazù - Argentina.

- 1995, 3.44 A. PELOSI, R. LOMBARDI
Fondi pensione: equilibrio economico-finanziario delle imprese.
- 1995, 3.45 Emanuele SALSANO, Domenico IANNONE
Economia e struttura produttiva nel salernitano dal secondo dopoguerra ad oggi.
- 1995, 3.46 Michele LA ROCCA
Empirical likelihood and linear combinations of functions of order statistics.
- 1995, 3.47 Michele LA ROCCA
L'uso del bootstrap nella verosimiglianza empirica.
- 1996, 3.48 Domenico RANESI
Le politiche CEE per lo sviluppo dei sistemi locali: esame delle diverse tipologie di intervento e tentativo di specificazione tassonomica.
- 1996, 3.49 Michele LA ROCCA
L'uso della verosimiglianza empirica per il confronto di due parametri di posizione.
- 1996, 3.50 Massimo SPAGNOLO
La domanda dei prodotti della pesca in Italia.
- 1996, 3.51 Cesare IMBRIANI, Filippo REGANATI
Macroeconomic stability and economic integration. The case of Italy.
- 1996, 3.52 Annarita GERMANI
Gli effetti della mobilitazione della riserva obbligatoria. Analisi sull'efficienza del suo utilizzo.
- 1996, 3.53 Massimo SPAGNOLO
A model of fish price formation in the north sea and the Mediterranean.
- 1996, 3.54 Fernanda MAZZOTTA
RTFL: problemi e soluzioni per i dati Panel.
- 1996, 3.55 Angela SPAGNUOLO
Concentrazione industriale e dimensione del mercato: il ruolo della spesa per pubblicità e R&D.
- 1996, 3.56 Giuseppina AUTIERO
The economic case for social norms.
- 1996, 3.57 Francesco GIORDANO
Sulla convergenza degli stimatori Kernel.
- 1996, 3.58 Tullio JAPPELLI, Marco PAGANO
The determinants of saving: lessons from Italy.

- 1997, 3.59 Tullio JAPPELLI
The age-wealth profile and the life-cycle hypothesis: a cohort analysis with a time series of cross sections of Italian households.
- 1997, 3.60 Marco Antonio MONACO
La gestione dei servizi di pubblico interesse.
- 1997, 3.61 Marcella ANZOLIN
L'albero della qualità dei servizi pubblici locali in Italia: metodologie e risultati conseguiti.
- 1997, 3.62 Cesare IMBRIANI, Antonio LOPES
Intermediazione finanziaria e sistema produttivo in un'area dualistica. Uno studio di caso.
- 1997, 3.63 Tullio JAPPELLI
Risparmio e liberalizzazione finanziaria nell'Unione europea.
- 1997, 3.64 Alessandra AMENDOLA
Analisi dei dati di sopravvivenza.
- 1997, 3.65 Francesco GIORDANO, Cira PERNA
Gli stimatori Kernel per la stima non parametrica della funzione di regressione.
- 1997, 3.66 Biagio DI SALVIA
*Le relazioni marittimo-commerciali nell'imperiale regio litorale austriaco nella prima metà dell'800.
I. Una riclassificazione delle Tafeln zur Statistik der Österreichischen Monarchie.*
- 1997, 3.67 Alessandra AMENDOLA
Modelli non lineari di seconda e terza generazione: aspetti teorici ed evidenze empiriche.
- 1998, 3.68 Vania SENA
L'analisi econometrica dell'efficienza tecnica. Un'applicazione agli ospedali italiani di zona.
- 1998, 3.69 Domenico CERBONE
Investimenti irreversibili.
- 1998, 3.70 Antonio GAROFALO
La riduzione dell'orario di lavoro è una soluzione al problema disoccupazione: un tentativo di analisi empirica.
- 1998, 3.71 Jacqueline MORGAN, Roberto RAUCCI
New convergence results for Nash equilibria.

- 1998, 3.72 Rosa FERRENTINO
Niels Henrik Abel e le equazioni algebriche.
- 1998, 3.73 Marco MICOCCI, Rosa FERRENTINO
Un approccio markoviano al problema della valutazione delle opzioni.
- 1998, 3.74 Rosa FERRENTINO, Ciro CALABRESE
Rango di una matrice di dimensione K .
- 1999, 3.75 Patrizia RIGANTI
L'uso della valutazione contingente per la gestione del patrimonio culturale: limiti e potenzialità.
- 1999, 3.76 Annamaria NESE
Il problema dell'inefficienza nel settore dei musei: tecniche di valutazione.
- 1999, 3.77 Gianluigi COPPOLA
Disoccupazione e mercato del lavoro: un'analisi su dati provinciali.
- 1999, 3.78 Alessandra AMENDOLA
Un modello soglia con eteroschedasticità condizionata per tassi di cambio.
- 1999, 3.79 Rosa FERRENTINO
Su un'applicazione della trasformata di Laplace al calcolo della funzione asintotica di non rovina.
- 1999, 3.80 Rosa FERRENTINO
Un'applicazione della trasformata di Laplace nel caso di una distribuzione di Erlang.
- 1999, 3.81 Angela SPAGNUOLO
Efficienza e struttura degli incentivi nell'azienda pubblica: il caso dell'industria sanitaria.
- 1999, 3.82 Antonio GAROFALO, Cesare IMBRIANI, Concetto Paolo VINCI
Youth unemployment: an insider-outsider dynamic approach.
- 1999, 3.83 Rosa FERRENTINO
Un modello per la determinazione del tasso di riequilibrio in un progetto di fusione tra banche.
- 1999, 3.84 DE STEFANIS, PORZIO
Assessing models in frontier analysis through dynamic graphics.
- 1999, 3.85 Annunziato GESUALDI
Inflazione e analisi delle politiche fiscali nell'U.E..
- 1999, 3.86 R. RAUCCI, L. TADDEO
Dalle equazioni differenziali alle funzioni e^x , $\log x$, a^x , $\log_a x$, x^x .

- 1999, 3.87 Rosa FERRENTINO
Sulla determinazione di numeri aleatori generati da equazioni algebriche.
- 1999, 3.88 C. PALMISANI, R. RAUCCI
Sulle funzioni circolari: una presentazione non classica.
- 2000, 3.89 Giuseppe STORTI, Pierluigi FURCOLO, Paolo VILLANI
A dynamic generalized linear model for precipitation forecasting.
- 2000, 3.90 Rosa FERRENTINO
Un procedimento risolutivo per l'equazione di Dickson.
- 2000, 3.91 Rosa FERRENTINO
Un'applicazione della mistura di esponenziali alla teoria del rischio.
- 2000, 3.92 Francesco GIORDANO, Michele LA ROCCA, Cira PERNA
Bootstrap variance estimates for neural networks regression models.
- 2000, 3.93 Alessandra AMENDOLA, Giuseppe STORTI
A non-linear time series approach to modelling asymmetry in stock market indexes.
- 2000, 3.94 Rosa FERRENTINO
Sopra un'osservazione di De Vylder.
- 2000, 3.95 Massimo SALZANO
Reti neurali ed efficacia dell'intervento pubblico: previsioni dell'inquinamento da traffico nell'area di Villa S. Giovanni.
- 2000, 3.96 Angela SPAGNUOLO
Concorrenza e deregolamentazione nel mercato del trasporto aereo in Italia.
- 2000, 3.97 Roberto RAUCCI, Luigi TADDEO
Teoremi ingannevoli.
- 2000, 3.98 Francesco GIORDANO
Una procedura per l'inizializzazione dei pesi delle reti neurali per l'analisi del trend.
- 2001, 3.99 Angela D'ELIA
Some methodological issues on multivariate modelling of rank data.
- 2001, 3.100 Roberto RAUCCI, Luigi TADDEO
Nuove classi di funzioni scalari quasiconcave generalizzate: caratterizzazioni ed applicazioni a problemi di ottimizzazione.
- 2001, 3.101 Adriana BARONE, Annamaria NESE
Some insights into night work in Italy.
- 2001, 3.102 Alessandra AMENDOLA, Marcella NIGLIO

Predictive distributions of nonlinear time series models.

- 2001, 3.103 Roberto RAUCCI
Sul concetto di certo equivalente nella teoria HSSB.
- 2001, 3.104 Roberto RAUCCI, Luigi TADDEO
On stackelberg games: a result of unicity.
- 2001, 3.105 Roberto RAUCCI
Una definizione generale e flessibile di insieme limitato superiormente in \mathfrak{R}^n
- 2001, 3.106 Roberto RAUCCI
Stretta quasiconcavit  nelle forme funzionali flessibili.
- 2001, 3.107 Roberto RAUCCI
Sugli insiemi limitati in \mathfrak{R}^m rispetto ai coni.
- 2001, 3.108 Roberto RAUCCI
Monotonie, isotonie e indecomponibilit  deboli per funzioni a valori vettoriali con applicazioni.
- 2001, 3.109 Roberto RAUCCI
Generalizzazioni del concetto di debole Kuhn-Tucker punto-sella.
- 2001, 3.110 Antonia Rosa GURRIERI, Marilene LORIZIO
Le determinanti dell'efficienza nel settore sanitario. Uno studio applicato.
- 2001, 3.111 Gianluigi COPPOLA
Studio di una provincia meridionale attraverso un'analisi dei sistemi locali del lavoro. Il caso di Salerno.
- 2001, 3.112 Francesco GIORDANO
Reti neurali per l'analisi del trend: un approccio per identificare la topologia della rete.
- 2001, 3.113 Marcella NIGLIO
Nonlinear time series models with switching structure: a comparison of their forecast performances.
- 2001, 3.114 Damiano FIORILLO
Capitale sociale e crescita economica. Review dei concetti e dell'evidenza empirica.
- 2001, 3.115 Roberto RAUCCI, Luigi TADDEO
Generalizzazione del concetto di continuit  e di derivabilit .
- 2001, 3.116 Marcella NIGLIO
Ricostruzione dei dati mancanti in serie storiche climatiche.

- 2001, 3.117 Vincenzo VECCHIONE
Mutamenti del sistema creditizio in un'area periferica.
- 2002, 3.118 Francesco GIORDANO, Michele LA ROCCA, Cira PERNA
Bootstrap variable selection in neural network regression models.
- 2002, 3.119 Roberto RAUCCI, Luigi TADDEO
Insiemi debolmente convessi e concavità in senso generale.
- 2002, 3.120 Vincenzo VECCHIONE
Know how locali e percorsi di sviluppo in aree e settori marginali.
- 2002, 3.121 Michele LA ROCCA, Cira PERNA
Neural networks with dependent data.
- 2002, 3.122 Pietro SENESI
Economic dynamics: theory and policy. A stability analysis approach.
- 2002, 3.123 Gianluigi COPPOLA
Stima di un indicatore di pressione ambientale: un'applicazione ai comuni della Campania.
- 2002, 3.124 Roberto RAUCCI
Sull'esistenza di autovalori e autovettori positivi anche nel caso non lineare.
- 2002, 3.125 Maria Carmela MICCOLI
Identikit di giovani lucani.
- 2002, 3.126 Sergio DESTEFANIS, Giuseppe STORTI
Convexity, productivity change and the economic performance of countries.
- 2002, 3.127 Giovanni C. PORZIO, Maria Prosperina VITALE
Esplorare la non linearità nei modelli Path.
- 2002, 3.128 Rosa FERRENTINO
Sulla funzione di Seal.
- 2003, 3.129 Michele LA ROCCA, Cira PERNA
Identificazione del livello intermedio nelle reti neurali di tipo feedforward.
- 2003, 3.130 Alessandra AMENDOLA, Marcella NIGLIO, Cosimo VITALE
The exact multi-step ahead predictor of SETARMA models.
- 2003, 3.131 Mariangela BONASIA
La dimensione ottimale di un sistema pensionistico: means tested vs programma universale.
- 2003, 3.132 Annamaria NESE
Abitazione e famiglie a basso reddito.

- 2003, 3.133 Maria Lucia PARRELLA
Le proprietà asintotiche del Local Polynomial Bootstrap.
- 2003, 3.134 Silvio GIOVE, Maurizio NORDIO, Stefano SILVONI
Stima della prevalenza dell'insufficienza renale cronica con reti bayesiane: analisi costo efficacia delle strategie di prevenzione secondaria.
- 2003, 3.135 Massimo SALZANO
Globalization, complexity and the holism of the italian school of public finance.
- 2003, 3.136 Giuseppina AUTIERO
Labour market institutional systems and unemployment performance in some Oecd countries.
- 2003, 3.137 Marisa FAGGINI
Recurrence analysis for detecting non-stationarity and chaos in economic times series.
- 2003, 3.138 Marisa FAGGINI, Massimo SALZANO
The reverse engineering of economic systems. Tools and methodology.
- 2003, 3.139 Rosa FERRENTINO
In corso di pubblicazione.
- 2003, 3.140 Rosa FERRENTINO, Roberto RAUCCI
Sui problemi di ottimizzazione in giochi di Stackelberg ed applicazioni in modelli economici.
- 2003, 3.141 Carmine SICA
In corso di pubblicazione.
- 2004, 3.142 Sergio DESTEFANIS, Antonella TADDEO, Maurizio TORNATORE
The stock of human capital in the Italian regions.
- 2004, 3.143 Elena Laureana DEL MERCATO
Edgeworth equilibria with private provision of public good.
- 2004, 3.144 Elena Laureana DEL MERCATO
Externalities on consumption sets in general equilibrium.
- 2004, 3.145 Rosa FERRENTINO, Roberto RAUCCI
Su alcuni criteri delle serie a termini non negativi.
- 2004, 3.146 Rosa FERRENTINO, Roberto RAUCCI
Legame tra le soluzioni di Minty e di Stempacenhia nelle disequazioni variazionali.

- 2004, 3.147 Gianluigi COPPOLA
In corso di pubblicazione.
- 2004, 3.148 Massimo Spagnolo
The Importance of Economic Incentives in Fisheries Management
- 2004, 3.149 F. Salsano
La politica monetaria in presenza di non perfetta osservabilità degli obiettivi del banchiere centrale.
- 2004, 3.150 A. Vita
La dinamica del cambiamento nella rappresentazione del territorio. Una mappa per i luoghi della Valle dell'Irno.
- 2004, 3.151 Celi
Empirical Explanation of vertical and horizontal intra-industry trade in the UK: a comment.
- 2004, 3.152 Amendola – P. Vitale
Self-Assessment and Career Choices: An On-line resource for the University of Salerno.
- 2004, 3.153 A. Amendola – R. Troisi
Introduzione all'economia politica dell'organizzazione: nozioni ed applicazioni.
- 2004, 3.154 A. Amendola – R. Troisi
Strumenti d'incentivo e modelli di gestione del personale volontario nelle organizzazioni non profit.
- 2004, 3.155 Lavinia Parisi
La gestione del personale nelle imprese manifatturiere della provincia di Salerno.
- 2004, 3.156 Angela Spagnuolo – Silvia Keller
La rete di accesso all'ultimo miglio: una valutazione sulle tecnologie alternative.
- 2005, 3.157 Davide Cantarelli
Elasticities of Complementarity and Substitution in Some Functional Forms. A Comparative Review.
- 2005, 3.158 Pietro Coretto – Giuseppe Storti
Subjective Expectations in Economics: a Statistical overview of the main findings.
- 2005, 3.159 Pietro Coretto – Giuseppe Storti
Moments based inference in small samples.

- 2005, 3.160 Massimo Salzano
Una simulazione neo-keynesiana ad agenti eterogeni.
- 2005, 3.161 Rosa Ferrentino
Su alcuni paradossi della teoria degli insiemi.
- 2005, 3.162 Damiano Fiorillo
Capitale sociale: uno o molti? Pochi.
- 2005, 3.163 Damiano Fiorillo
Il capitale sociale conta per outcomes (macro) economici?.
- 2005, 3.164 Damiano Fiorillo – Guadalupi Luigi
*Attività economiche nel distretto industriale di Nocera inferiore – Gragnano.
Un'analisi su Dati Tagliacarne.*
- 2005, 3.165 Rosa Ferrentino
Pointwise well-posedness in vector optimization and variational inequalities.
- 2005, 3.166 Roberto Iorio
La ricerca universitaria verso il mercato per il trasferimento tecnologico e rischi per l'Open Science: posizioni teoriche e filoni di indagine empirica.
- 2005, 3.167 Marisa Faggini
The chaotic system and new perspectives for economics methodology. A note.
- 2005, 3.168 Francesco Giordano
Weak consistent moving block bootstrap estimator of sampling distribution of CLS estimators in a class of bilinear models
- 2005, 3.169 Edgardo Sica
Tourism as determinant of economic growth: the case of south-east asian countries.
- 2005, 3.170 Rosa Ferrentino
On Minty variational inequalities and increasing along rays functions.
- 2005, 3.171 Rosa Ferrentino
On the Minty and Stampacchia scalar variational inequalities
- 2005, 3.172 Destefanis - Storti
A procedure for detecting outliers in frontier estimation
- 2005, 3.173 Destefanis - Storti
Evaluating business incentives through DEA. An analysis on capitalia firm data

- 2005, 3.174 Nese – O'Higgins
In and out of the capitalia sample: evaluating attrition bias.
- 2005, 3.175 Maria Patrizia Vittoria
Il Processo di terziarizzazione in Campania. Analisi degli indicatori principali nel periodo 1981-2001
- 2005, 3.176 Sergio Destefanis – Giuseppe Mastromatteo
Inequality and labour-market performance. A survey beyond an elusive trade-off.
- 2006, 3.177 Giuseppe Storti
Modelling asymmetric volatility dynamics by multivariate BL-GARCH models
- 2006, 3.178 Lucio Valerio Spagnolo – Mario Cerrato
No euro please, We're British!
- 2006, 3.179 Maria Carmela Miccoli
Invecchiamento e seconda transizione demografica
- 2006, 3.180 Maria Carmela Miccoli – Antonio Cortese
Le scuole italiane all'estero: una realtà poco nota
- 2007, 3.181 Rosa Ferrentino
Variational inequalities and optimization problems
- 2007, 3.182 Lavinia Parisi
Estimating capability as a latent variable: A Multiple Indicators and Multiple Causes Approach. The example of health
- 2007, 3.183 Rosa Ferrentino
Well-posedness, a short survey
- 2007, 3.184 Roberto Iorio – Sandrine Labory – Daniele Paci
Relazioni tra imprese e università nel biotech-salute dell'Emilia Romagna. Una valutazione sulla base della co-authorship delle pubblicazioni scientifiche
- 2007, 3.185 Lavinia Parisi
Youth Poverty after leaving parental home: does parental income matter?

Stampa a cura della C.U.S.L. Cooperativa Universitaria Studio e Lavoro, Via Ponte Don Melillo, Fisciano per conto Del Dipartimento di Scienze Economiche e Statistiche
Finito di stampare il 15 maggio 2007